arXiv:1701.01905v1 [math.AG] 8 Jan 2017

KORFF F-SIGNATURES OF HIRZEBRUCH SURFACES

DAISUKE HIROSE AND TADAKAZU SAWADA

Abstract. M. Von Korff introduced the F-signature of a normal projective variety, and computed the F-signature of the product of two projective lines $\mathbb{P}^1 \times \mathbb{P}^1$. In this paper, we compute F-signatures of Hirzebruch surfaces.

Introduction

Let R be a d-dimensional Noetherian local ring of prime characteristic p with perfect residue field. Let $F : R \to R$ be the Frobenius map, that is, $F(x) = x^p$ for all elements x of R. We denote by $F_*^e R$ the R-module whose abelian group structure is inherited by R, and whose R-module structure is given by the e-th iterated Frobenius map F^e. If F_*R is finitely generated as an R-module, we say that R is *F-finite*. Let the R-module $F_*^e R$ be decomposed as $F_*^e R \cong R^{\oplus a_e} \oplus M_e$, where M_e does not have free summands. In [HL02], C. Huneke and G. Leuschke introduced the notion of the F-signature $s(R)$ of R as

$$s(R) = \lim_{e\to\infty} \frac{a_e}{p^{ed}}.$$

The existence of the limit $\lim_{e\to\infty}(a_e/p^{ed})$ is not trivial. In [Tuc12], K. Tucker proved that the limit exists (under the assumption that R is F-finite). The notion of the F-signature is generalized by M. Hashimoto and Y. Nakajima in [HN15], and A. Sannai in [San15]. As a global analogue, M. Von Korff introduced the F-signature of a normal projective variety, which we call the Korff F-signature, in [Kor12]. He computed the Korff F-signature of the product of two projective lines $\mathbb{P}^1 \times \mathbb{P}^1$. In this paper, we compute Korff F-signatures of Hirzebruch surfaces.

In Section 1, we review generalities on Korff F-signatures. In Section 2, we compute Korff F-signatures of projective spaces and Hirzebruch surfaces. In Section 3 and 4, we gather together the figures and the source code of wxMaxima in the computation of F-signatures of Hirzebruch surfaces.

1. Preliminaries

In this section, we review generalities on Korff F-signatures. See [Kor12] for details.

Let k be a field of positive characteristic p. Let X be a normal projective variety over k, and let D be a $\mathbb{Q}$-divisor on X. We denote by $\mathrm{Sec}(X,D)$ the section ring $\bigoplus_{n\geq 0}\Gamma(X,\mathcal{O}_X(nD))$ of D. Suppose that $\mathrm{Sec}(X,D)$ is a finitely generated k-algebra of dimension at least two. If $c\in\mathbb{N}$ is sufficiently divisible, then there exists an ample line bundle L on X such that $\mathrm{Sec}(X,cD)$ is isomorphic to a normal section ring $\mathrm{Sec}(X,L)=\bigoplus_{n\geq 0}\Gamma(X,L^{\otimes n})$ of L. We assume that c is sufficiently divisible, and define the *Korff F-signature $s(X,D)$ of X along D* to be $c\cdot s(\mathrm{Sec}(X,cD))$. (The F-signature of a graded ring is defined in the same way as the case of local rings.)

Let $N\cong\mathbb{Z}^n$ be a lattice, and let M be the dual lattice of N. We denote $N\otimes_{\mathbb{Z}}\mathbb{R}$ (resp. $M\otimes_{\mathbb{Z}}\mathbb{R}$) by $N_{\mathbb{R}}$ (resp. $M_{\mathbb{R}}$). Let Σ be a fan in $N_{\mathbb{R}}$. We denote by X_Σ the toric variety corresponding to Σ.

Let Σ be a complete fan in $N_{\mathbb{R}}$ with the primitive generators $v_1,\dots,v_n$. Let $D_1,\dots,D_n$ be the torus-invariant prime Weil divisors corresponding to $v_1,\dots,v_n$, respectively. Let $D=\sum_{i=1}^n a_iD_i$ be a torus-invariant Weil divisor on X. The *polytope associated to the divisor D* is defined to be the polytope $\{u\in M_{\mathbb{R}}|u\cdot v_i\geq -a_i \text{ for all } i\}$, and denoted by P_D. We may assume that P_D contains the origin. Then we denote by M_D the lattice $M\cap(P_D\cdot\mathbb{R})$, where $P_D\cdot\mathbb{R}$ is the $\mathbb{R}$-vector subspace spanned by P_D in $M_{\mathbb{R}}$. We denote by N_D the dual lattice of M_D. We define I_D to be the subset of $\{1,\dots,n\}$ such that the hyperplane $\{u|u\cdot v_i=-a_i\}$ determines a facet of P_D for $i\in I_D$, and $P_{\Sigma,D}$ to be the polytope $\{(v,t)\in(M_D\times\mathbb{Z})_{\mathbb{R}}|0\leq(v,t)\cdot(v_i,a_i)<1 \text{ for all } i\in I_D\}$.

Suppose that P_D is not full-dimensional. Let $\pi:N\to N_D$ be the natural projection map. Let c_i be the rational number such that $\pi(c_iv_i)$ is the primitive generator for its ray in $(N_D)_{\mathbb{R}}$. We define $P'_{\Sigma,D}$ to be the polytope $\{(v,t)\in(M_D\times\mathbb{Z})_{\mathbb{R}}|0\leq(v,t)\cdot c_i(v_i,a_i)<1 \text{ for all } i\in I_D\}$.

Theorem 1.1 ([Kor12], Proposition 4.4.4, Corollary 4.4.7)**.** *Let X be a d-dimensional projective toric variety over k, and let Σ be the fan corresponding to X. Let $v_1,\dots,v_n$ be the primitive generators of Σ, and let $D_1,\dots,D_n$ be the torus-invariant prime Weil divisors corresponding to $v_1,\dots,v_n$, respectively. Let $D=\sum_{i=1}^n a_iD_i$ be a (non-zero) effective torus-invariant Weil divisor on X. Then:*
(1) *If P_D is full-dimensional, then $s(X,D)=\mathrm{Vol}(P_{\Sigma,D})$.*
(2) *If P_D is not full-dimensional, then $s(X,D)=\mathrm{Vol}(P'_{\Sigma,D})$.*

2. Examples

M. Von Korff computed Korff F-signatures of some toric varieties including the product of two projective lines $\mathbb{P}^1 \times \mathbb{P}^1$ in [Kor12]. In this section, we compute Korff F-signatures of projective spaces and Hirzebruch surfaces.

Example 2.1 (Projective spaces $\mathbb{P}^n$)**.** Let $N = \mathbb{Z}^n$ with standard basis $e_1, \ldots, e_n$. Let $e_0 = -\sum_{i=1}^n e_i$, and let Σ be the fan in $N_{\mathbb{R}}$ consisting of the cones generated by all proper subsets of $\{e_0, e_1, \ldots, e_n\}$. We see that X_Σ is isomorphic to the n-dimensional projective space $\mathbb{P}^n$. Let D_i be the torus-invariant prime Weil divisor corresponding to the ray $\mathbb{R}_+ e_i$ in $N_{\mathbb{R}}$ for $0 \leq i \leq n$.

Let $D = \sum_{i=0}^n a_i D_i$ be a (non-zero) effective torus-invariant Weil divisor. We see that

$$P_D = \left\{ (x_1, \ldots, x_n) \in M_{\mathbb{R}} \;\middle|\; \begin{array}{l} \sum_{i=1}^n x_i \leq a_0, \\ x_1 \geq -a_1, \\ \cdots \\ x_n \geq -a_n \end{array} \right\}.$$

Since $a_i \neq 0$ for some i, P_D has an interior point, and the equations $\sum_{i=1}^n x_i = a_0, x_1 = -a_1, \ldots, x_n = -a_n$ define the facets of P_D, respectively. Hence we have $I_D = \{0, 1, \ldots, n\}$. In what follows, we compute the volume of the n-dimensional parallelepipeds

$$P_{\Sigma,D} = \left\{ (x_1, \ldots, x_n, t) \in (M_D \times \mathbb{Z})_{\mathbb{R}} \;\middle|\; \begin{array}{l} 0 \leq -\sum_{i=1}^n x_i + a_0 t < 1, \\ 0 \leq x_1 + a_1 t < 1, \\ \cdots \\ 0 \leq x_n + a_n t < 1 \end{array} \right\}.$$

Let $a = \sum_{i=0}^n a_i$. The point of the intersection of the $n+1$ hyperplanes defined by $-\sum_{i=1}^n x_i + a_0 t = 1, x_1 + a_1 t = 0, \ldots, x_n + a_n t = 0$ is $(-a_1/a, \ldots, -a_n/a, 1/a)$. This is given by the following elementary transformations of matrices:

$$\left(\begin{array}{ccccc|c} -1 & -1 & \cdots & -1 & a_0 & 1 \\ 1 & 0 & \cdots & 0 & a_1 & 0 \\ & & \cdots & & & \\ 0 & 0 & \cdots & 1 & a_n & 0 \end{array}\right) \to \left(\begin{array}{ccccc|c} 0 & 0 & \cdots & 0 & a & 1 \\ 1 & 0 & \cdots & 0 & a_1 & 0 \\ & & \cdots & & & \\ 0 & 0 & \cdots & 1 & a_n & 0 \end{array}\right)$$

$$\to \left(\begin{array}{ccccc|c} 0 & 0 & \cdots & 0 & a & 1 \\ 1 & 0 & \cdots & 0 & 0 & -a_1/a \\ & & \cdots & & & \\ 0 & 0 & \cdots & 1 & 0 & -a_n/a \end{array}\right) \to \left(\begin{array}{ccccc|c} 1 & 0 & \cdots & 0 & 0 & -a_1/a \\ & & \cdots & & & \\ 0 & 0 & \cdots & 1 & 0 & -a_n/a \\ 0 & 0 & \cdots & 0 & 1 & 1/a \end{array}\right).$$

By the same argument, we see that the point of the intersection of the $n+1$ hyperplanes defined by $-\sum_{i=1}^{n} x_i + a_0 t = 0$, $x_l + a_l t = 0$ $(l \neq k)$, $x_k + a_k t = 1$ is $(-a_1/a, \ldots, 1-a_k/a, \ldots, -a_n/a, 1/a)$ for each $1 \leq k \leq n$.

Let $v_0 = {}^t(-a_1/a, \ldots, -a_n/a, 1/a)$, and

$$v_k = {}^t(-a_1/a, \ldots, 1 - a_k/a, \ldots, -a_n/a, 1/a)$$

for $1 \leq k \leq n$. Since $P_{\Sigma,D}$ is spanned by $v_0, v_1, \ldots, v_n$, we have

$$\begin{aligned}
s(\mathbb{P}^n, D) &= \mathrm{Vol}(P_{\Sigma,D}) \\
&= |\det(v_0, v_1, \ldots, v_n)| \\
&= \left| \det \begin{pmatrix} -a_1/a & 1-a_1/a & -a_1/a & \cdots & -a_1/a \\ -a_2/a & -a_2/a & 1-a_2/a & \cdots & -a_2/a \\ & & & \cdots & \\ -a_n/a & -a_n/a & -a_n/a & \cdots & 1-a_n/a \\ 1/a & 1/a & 1/a & \cdots & 1/a \end{pmatrix} \right| \\
&= 1/a^{n+1} \left| \det \begin{pmatrix} -a_1 & a-a_1 & -a_1 & \cdots & -a_1 \\ -a_2 & -a_2 & a-a_2 & \cdots & -a_2 \\ & & & \cdots & \\ -a_n & -a_n & -a_n & \cdots & a-a_n \\ 1 & 1 & 1 & \cdots & 1 \end{pmatrix} \right| \\
&= 1/a^{n+1} \left| \det \begin{pmatrix} 0 & a & 0 & \cdots & 0 \\ 0 & 0 & a & \cdots & 0 \\ & & & \cdots & \\ 0 & 0 & 0 & \cdots & a \\ 1 & 1 & 1 & \cdots & 1 \end{pmatrix} \right| \\
&= 1/a = 1/(a_0 + a_1 + \cdots + a_n).
\end{aligned}$$

The following computation is the main result.

Example 2.2 (Hirzebruch surfaces)**.** All figures of this example are in Section 3. Let r be a positive integer, and let $N = \mathbb{Z}^2$ with standard basis e_1 and e_2. Let $v_1 = e_1, v_2 = e_2, v_3 = -e_2$, and $v_4 = -e_1 + re_2$. We consider a fan Σ_r in $N_{\mathbb{R}} = \mathbb{R}^2$. The fan Σ_r consists the four two-dimensional cones $\langle v_1, v_2\rangle, \langle v_1, v_3\rangle, \langle v_2, v_4\rangle$, and $\langle v_3, v_4\rangle$, and all their faces. The Hirzebruch surface $X = X_{\Sigma_r}$ is defined by the fan Σ_r (see, e.g., [CLS11], Example 3.1.16.). Let D_i be the torus-invariant prime divisors of X corresponding to v_i. We have linear equivalents $D_1 \sim D_4$ and $D_3 \sim D_2 + rD_4$ in the divisor class group of X. For an effective $\mathbb{Q}$-divisor $D = \sum_{i=1}^{4} a_i D_i$, we have

$$\begin{aligned}
D \sim &\quad a_1 D_1 + (a_2 + a_3) D_2 + (ra_3 + a_4) D_4 \\
\sim &\quad (a_1 + ra_3 + a_4) D_1 + (a_2 + a_3) D_2.
\end{aligned}$$

Hence $\mathcal{O}(D) \cong \mathcal{O}((a_1 + ra_3 + a_4)D_1 + (a_2 + a_3)D_2)$. Without loss of generality, we only compute the F-signature $s(X, D)$ of X with respect to a (non-zero) effective $\mathbb{Q}$-divisor $D = a_1D_1 + a_2D_2$.

The polytope P_D is to be

$$P_D = \left\{ (x, y) \in M_{\mathbb{R}} \;\middle|\; \begin{array}{l} x \geq -a_1,\ y \geq -a_2 \\ y \leq 0,\ -x + ry \geq 0 \end{array} \right\}.$$

By the figure 1, we see that P_D is a square if and only if $ra_2 < a_1$. In that case, $I_D = \{1, 2, 3, 4\}$. On the other hand, if $ra_2 \geq a_1$, then P_D is a triangle. That implies $I_D = \{1, 3, 4\}$.
(1) Suppose that P_D is a square, i.e., $ra_2 < a_1$. Then $I_D = \{1, 2, 3, 4\}$ Therefore

$$P_{\Sigma,D} = \left\{ (x, y, t) \in (M_D \times \mathbb{Z})_{\mathbb{R}} \;\middle|\; \begin{array}{l} 0 \leq x + a_1t < 1, \\ 0 \leq y + a_2t < 1, \\ 0 \leq -y < 1, \\ 0 \leq -x + ry < 1 \end{array} \right\}.$$

We define polygons Q, $R(x)$, and $S(x)$ to be

$$\begin{aligned} Q &= \left\{ (y, t) \in \mathbb{R}^2 \;\middle|\; \begin{array}{l} 0 \leq y + a_2t < 1, \\ 0 \leq -y < 1 \end{array} \right\} \\ &= \left\{ (y, t) \in \mathbb{R}^2 \;\middle|\; \begin{array}{l} -a_2t \leq y < 1 - a_2t, \\ -1 < y \leq 0 \end{array} \right\}, \\ R(x) &= \left\{ (y, t) \in \mathbb{R}^2 \;\middle|\; \begin{array}{l} 0 \leq x + a_1t < 1, \\ 0 \leq -x + ry < 1 \end{array} \right\} \\ &= \left\{ (y, t) \in \mathbb{R}^2 \;\middle|\; \begin{array}{l} -x/a_1 \leq t < -x/a_1 + 1/a_1, \\ x/r \leq y < x/r + 1/r \end{array} \right\}, \end{aligned}$$

and $S(x) = Q \cap R(x)$.(See the figure 2.)

Then Q is a parallelogram, $R(x)$ is a rectangle, and

$$P_{\Sigma,D} = \{(x, y, t) \in (M_D \times \mathbb{Z})_{\mathbb{R}} | x_{\min} \leq x \leq x_{\max}, (y, t) \in S(x)\},$$

where $x_{\min} = \min\{x \in \mathbb{R} | S(x) \neq \emptyset\}$ and $x_{\max} = \max\{x \in \mathbb{R} | S(x) \neq \emptyset\}$. Therefore

$$s(X, D) = \mathrm{Vol}(P_{\Sigma,D}) = \int_{x_{\min}}^{x_{\max}} \mathrm{Area}(S(x))\, dx.$$

We define some points and lines in the ty-plane appearing in our argument. Let A, B, C, and D be the vertices of the rectangle $R(x)$.

In particular,

$$A = \left(-\frac{1}{a_1}x, \frac{1}{r}x\right), B = \left(-\frac{1}{a_1}x + \frac{1}{a_1}, \frac{1}{r}x\right),$$
$$C = \left(-\frac{1}{a_1}x + \frac{1}{a_1}, \frac{1}{r}x + \frac{1}{r}\right), \text{and } D = \left(-\frac{1}{a_1}x, \frac{1}{r}x + \frac{1}{r}\right).$$

The point A moves along the line $y = -a_1 t/r$, denoted by l_1. The points B and D move along the line $y = -a_1 t/r + 1/r$, denoted by l_2. The point C moves along the line $y = -a_1 t/r + 2/r$, denoted by l_3.

We denote the vetices of the parallelogram Q except the origin O by E, F, and G. That is,

$$E = \left(\frac{1}{a_2}, -1\right), F = \left(\frac{2}{a_2}, -1\right), \text{and } G = \left(\frac{1}{a_2}, 0\right).$$

Let H and I be the intersection points of the t-axis with l_2 and l_3, respectively. We denote the intersection points of the line $y = -1$ with l_1, l_2, and l_3 by J, K, and L, respectively. That is,

$$H = \left(\frac{1}{a_1}, 0\right), \qquad I = \left(\frac{2}{a_1}, 0\right),$$
$$J = \left(\frac{r}{a_1}, -1\right), \quad K = \left(\frac{r+1}{a_1}, -1\right), \quad \text{and } L = \left(\frac{r+2}{a_1}, -1\right).$$

For the point A, we denote the t-coordinate of A and y-coordinate of A by A_t and A_y, respectively. We use the same notation for all points in the ty-plane. For example, $B_t = -x/a_1 + 1/a_1$ and $J_y = -1$.

The shape of the polygon $P_{\Sigma,D}$ depends on how the lines l_1, l_2 and l_3 across the parallelogram Q. We divide into five subcases from (1-1) through (1-5):

(1-1) We assume that $L_t \leq E_t$. That is, $(r+2)/a_1 \leq 1/a_2$. This is equivalent to $(r+2)a_2 \leq a_1$. Then $2a_2 \leq a_1$. This is equivalent to $2/a_1 \leq 1/a_2$. Hence $I_t \leq G_t$. (See the figure 4.)

We denote the intersection points of the line OE with l_2 and l_3 by T and U, respectively. Then

$$T = \left(\frac{1}{a_1 - ra_2}, -\frac{a_2}{a_1 - ra_2}\right), \text{ and } U = \left(\frac{2}{a_1 - ra_2}, -\frac{2a_2}{a_1 - ra_2}\right).$$

If $C = U$, then $(-x/a_1 + 1/a_1, x/r + 1/r) = (2/(a_1 - ra_2), -2a_2/(a_1 - ra_2))$. Hence $x = -(a_1 + ra_2)/(a_1 - ra_2)$. If $D = T$, then $(-x/a_1, x/r + 1/r) = (1/(a_1 - ra_2), -a_2/(a_1 - ra_2))$. Hence $x = -a_1/(a_1 - ra_2)$. If $B = T$, then $(-x/a_1 + 1/a_1, x/r) = (1/(a_1 - ra_2), -a_2/(a_1 - ra_2))$. Hence $x = -ra_2/(a_1 - ra_2)$. If $C = I$ and $D = H$, then $(-x/a_1 +$

$1/a_1, x/r + 1/r) = (2/a_1, 0)$. Hence $x = -1$. If $A = O$ and $B = H$, then $(-x/a_1, x/r) = (0, 0)$. Hence $x = 0$.

Since $0 < ra_2 < a_1$, we have $ra_2 < a_1 < a_1 + ra_2$ and $a_1 - ra_2 > 0$. Hence $-(a_1 + ra_2)/(a_1 - ra_2) < -a_1/(a_1 - ra_2) < -ra_2/(a_1 - ra_2)$. Since $a_1 - ra_2 < a_1$, we have $-a_1/(a_1 - ra_2) < -1$. If $2ra_2 \leq a_1$, then $-1 \leq -ra_2/(a_1 - ra_2)$. Therefore

$$-\frac{a_2 + ra_2}{a_1 - ra_2} < -\frac{a_1}{a_1 - ra_2} < -1 < -\frac{ra_2}{a_1 - ra_2} < 0.$$

On the other hand, if $2ra_2 > a_1$, then $-ra_2/(a_1 - ra_2) < -1$. Therefore

$$-\frac{a_2 + ra_2}{a_1 - ra_2} < -\frac{a_1}{a_1 - ra_2} < -\frac{ra_2}{a_1 - ra_2} < -1 < 0.$$

For various values of x, we consider shapes of $S(x)$ and areas of them.

If $-(a_2 + ra_2)/(a_1 - ra_2) < x < -a_1/(a_1 - ra_2)$, then $S(x)$ is a triangle as in the figure 5. Hence

$$\begin{aligned}\mathrm{Area}(S(x)) &= \frac{1}{2}\left\{\left(\frac{1}{r}x + \frac{1}{r}\right) - \left(\frac{a_2}{a_1}x - \frac{a_2}{a_1}\right)\right\} \cdot \left\{\left(-\frac{1}{a_1}x + \frac{1}{a_1}\right) - \left(-\frac{1}{ra_2}x - \frac{1}{ra_2}\right)\right\} \\ &= \frac{1}{2a_1^2 a_2 r^2}\left\{(a_1 - ra_2)x + (a_1 + ra_2)\right\}^2.\end{aligned}$$

(1-1-1) Suppose that $2ra_2 \leq a_1$.

If $-a_1/(a_1 - ra_2) < x < -1$, then $S(x)$ is a trapezoid as in the figure 6. Hence

$$\begin{aligned}\mathrm{Area}(S(x)) &= \frac{1}{2}\left\{\left(\frac{1}{r}x + \frac{1}{r} - \frac{a_2}{a_1}x\right) + \left(\frac{1}{r}x + \frac{1}{r} - \frac{a_2}{a_1}x + \frac{a_2}{a_1}\right)\right\} \cdot \frac{1}{a_1} \\ &= \frac{1}{2a_1^2 r}\left\{2(a_1 - ra_2)x + 2a_1 + ra_2\right\}.\end{aligned}$$

We denote this area by V_1.

If $-1 < x < -ra_2/(a_1 - ra_2)$, then $S(x)$ is a tropezoid as in the figure 7. Hence

$$\begin{aligned}\mathrm{Area}(S(x)) &= \frac{1}{2}\left(-\frac{a_2}{a_1}x - \frac{a_2}{a_1}x + \frac{a_2}{a_1}\right) \cdot \frac{1}{a_1} \\ &= \frac{a_2}{2a_1^2}(1 - 2x).\end{aligned}$$

We denote this area by V_2.

If $-ra_2/(a_1 - ra_2) < x < 0$, then $S(x)$ is a pentagon as in the figure 8. Hence

$$\begin{aligned}\mathrm{Area}(S(x)) &= -\frac{1}{r}x \cdot \frac{1}{a_1} - \frac{1}{2}\left(\frac{a_2}{a_1}x - \frac{1}{r}x\right) \cdot \left\{-\frac{1}{ra_2}x - \left(-\frac{1}{a_1}x\right)\right\} \\ &= -\frac{1}{ra_1}x - \frac{(a_1 - ra_2)^2}{2r^2a_1^2a_2}x^2.\end{aligned}$$

We denote this area by V_3.

Therefore

$$\begin{aligned}s(X, D) &= \mathrm{Vol}(P_{\Sigma,D}) \\ &= \int_{-\frac{a_2+ra_2}{a_1-ra_2}}^{-\frac{a_1}{a_1-ra_2}} \frac{1}{2a_1^2a_2r^2}\left\{(a_1 - ra_2)x + (a_1 + ra_2)\right\}^2 dx \\ &\quad + \int_{-\frac{a_1}{a_1-ra_2}}^{-1} V_1\, dx + \int_{-1}^{-\frac{ra_2}{a_1-ra_2}} V_2\, dx + \int_{-\frac{ra_2}{a_1-ra_2}}^{0} V_3\, dx \\ &= \frac{a_2}{a_1(a_1 - a_2r)}.\end{aligned}$$

In the above calculation, we use wxMaxima (see Section 4 (%o1)).

(1-1-2) Suppose that $2ra_2 > a_1$. If $-a_1/(a_1 - ra_2) < x < -ra_2/(a_1 - ra_2)$, then $S(x)$ is a trapezoid as in the figure 9, which is the same shape as that in the case where $-a_1/(a_1 - ra_2) < x < -1$ on (1-1-1). Then $\mathrm{Area}(S(x)) = V_1$.

If $-ra_2/(a_1 - ra_2) < x < -1$, then $S(x)$ is a pentagon as in the figure 10. Hence

$$\begin{aligned}\mathrm{Area}(S(x)) &= \left(\frac{1}{r}x + \frac{1}{r} - \frac{1}{r}\right) \cdot \frac{1}{a_1} - \frac{1}{2}\left(\frac{a_2}{a_1}x - \frac{1}{r}x\right) \cdot \left\{-\frac{1}{ra_2}x - \left(-\frac{1}{a_1}x\right)\right\} \\ &= \frac{1}{ra_1} - \frac{(a_1 - ra_2)^2}{2a_1^2a_2r^2}x^2.\end{aligned}$$

We denote this area by V_4.

If $-1 < x < 0$, then $S(x)$ is a pentagon as in the figure 11, which is the same shape as that in the case where $-ra_2/(a_1 - ra_2) < x < 0$ on (1-1-1). Then $\mathrm{Area}(S(x)) = V_3$.

Therefore

$$\begin{aligned} s(X,D) &= \mathrm{Vol}(P_{\Sigma,D}) \\ &= \int_{-\frac{a_2+ra_2}{a_1-ra_2}}^{-\frac{a_1}{a_1-ra_2}} \frac{1}{2a_1^2a_2r^2}\left\{(a_1-ra_2)x+(a_1+ra_2)\right\}^2 dx \\ &\quad + \int_{-\frac{a_1}{a_1-ra_2}}^{-\frac{ra_2}{a_1-ra_2}} V_1\,dx + \int_{-\frac{ra_2}{a_1-ra_2}}^{-1} V_4\,dx + \int_{-1}^{0} V_3\,dx \\ &= \frac{a_2}{a_1(a_1-a_2r)}, \end{aligned}$$

(see Section 4 (%o2)).
(1-2) We assume that $K_t < E_t < L_t$. That is, $(r+1)/a_1 < 1/a_2 < (r+2)/a_1$. This is equivalent to $(r+1)a_2 < a_1 < (r+2)a_2$. Then $2a_2 < a_1$. We have $I_t \leq G_t$. Since $(r+1)a_2 < a_1$ and $a_2 \leq ra_2 < a_1$, we have $(r+2)a_2 < 2a_1$. Therefore $L_t < F_t$. The point T denotes the same in the case (1-1). (See the figure 12.)

If $C = L$ and $D = K$, then $(-x/a_1 + 1/a_1, x/r + 1/r) = ((r+2)/a_1, -1)$. Hence $x = -(r+1)$. If $B_t = E_t$, then $-(1/a_1)x + 1/a_1 = 1/a_2$. Hence $x = -(a_1 - a_2)/a_2$. By the same argument in the case (1-1), we have the following four values of x. If $D = T$, then $x = -a_1/(a_1 - ra_2)$. If $B = T$, then $x = -ra_2/(a_1 - ra_2)$. If $C = I$ and $D = H$, then $x = -1$. If $A = O$ and $B = H$, then $x = 0$.

Since $(r+1)a_2 < a_1$, we have $r(r+1)a_2 + a_1 < (r+1)a_1$. Hence $a_1 < (r+1)(a_1 - ra_2)$. Therefore $-(r+1) < -a_1/(a_1 - ra_2)$. If $a_1/(a_1 - ra_2) > (a_1 - a_2)/a_2$, then $a_1^2 - (r+2)a_2a_1 + ra_2^2 > 0$. We have

$$\frac{(r+2) - \sqrt{r^2+4}}{2}a_2 < a_1 < \frac{(r+2) + \sqrt{r^2+4}}{2}a_2.$$

By the assumption that $(r+1)a_2 < a_1 < (r+2)a_2$, we have $a_1/(a_1 - ra_2) > (a_1 - a_2)/a_2$ in case (1-2). Therefore $-a_1/(a_1 - ra_2) < -(a_1 - a_2)/a_2$. Since $2a_2 < a_1$, we have $a_2 < a_1 - a_2$. Hence $-(a_1 - a_2)/a_2 < -1$. The assumption that $(r+1)a_2 < a_1$ gives the inequalities $ra_2 < a_1 - a_2$ and $a_2 < a_1 - ra_2$. Those imply $ra_2/(a_1 - ra_2) < (a_1 - a_2)/a_2$. Therefore $-(a_1 - a_2)/a_2 < -ra_2/(a_1 - ra_2)$. Finally, we compare the inequality between -1 and $-ra_2/(a_1 - ra_2)$. Suppose that $-1 < -ra_2/(a_1 - ra_2)$. Then $a_1 > 2ra_2$. With the assumption that $a_1 < (r+2)a_2$, we have $2ra_2 < (r+2)a_2$. Hence $r < 2$. That is, $r = 1$. Therefore if $r = 1$, then

$$-(r+1) < -\frac{a_1}{a_1 - ra_2} < -\frac{a_1 - a_2}{a_2} < -1 < -\frac{ra_2}{a_1 - ra_2} < 0.$$

By the same argument, if $-ra_2/(a_1-ra_2) < -1$, then $1 < r$. Therefore if $r \geq 2$, then

$$-(r+1) < -\frac{a_1}{a_1 - ra_2} < -\frac{a_1 - a_2}{a_2} < -\frac{ra_2}{a_1 - ra_2} < -1 < 0.$$

We remark that $-ra_2/(a_1 - ra_2) \neq -1$, because r is integer.

For various values of x, we consider shapes of $S(x)$ and areas of them.

If $-(r+1) < x < -a_1/(a_1 - ra_2)$, then $S(x)$ is a trapezoid as in the figure 13. Hence

$$\begin{aligned}
\mathrm{Area}(S(x)) &= \frac{1}{2}\left\{\left(-\frac{1}{a_1}x + \frac{1}{a_1} - \left(-\frac{1}{ra_2}x - \frac{1}{ra_2}\right)\right) + \left(-\frac{1}{a_1}x + \frac{1}{a_1} - \frac{1}{a_2}\right)\right\} \\
&\quad \times \left(\frac{1}{r}x + \frac{1}{r} - (-1)\right) \\
&= \frac{1}{2a_1a_2r^2}\left\{(a_1 - 2a_2r)x + 2a_2r + a_1 - a_1r\right\}(x + r + 1).
\end{aligned}$$

If $-a_1/(a_1 - ra_2) < x < -(a_1 - a_2)/a_2$, then $S(x)$ is a pentagon as in the figure 14. Hence

$$\begin{aligned}
\mathrm{Area}(S(x)) &= \left\{\frac{1}{r}x + \frac{1}{r} - (-1)\right\} \cdot \frac{1}{a_1} - \frac{1}{2}\left\{\frac{1}{a_2} - \left(-\frac{1}{a_1}x\right)\right\} \cdot \left\{\frac{a_2}{a_1}x - (-1)\right\} \\
&= \frac{1}{a_1r}(x + r + 1) - \frac{1}{2a_1^2a_2}(a_2x + a_1)^2.
\end{aligned}$$

We denote this area by V_5.

(1-2-1) Suppose that $r = 1$. If $-(a_1 - a_2)/a_2 < x < -1$, then $S(x)$ is a trapezoid as in the figure 15, which is the same shape as that in the case where $-a_1/(a_1 - ra_2) < x < -1$ on (1-1-1). Hence $\mathrm{Area}(S(x)) = V_1$.

If $-1 < x < -ra_2/(a_1 - ra_2)$, then $S(x)$ is a trapezoid as in the figure 16, which is the same shape as that in the case where $-1 < x < -ra_2/(a_1 - ra_2)$ on (1-1-1). Hence $\mathrm{Area}(S(x)) = V_2$.

If $-ra_2/(a_1 - ra_2) < x < 0$, then $S(x)$ is a pentagon as in the figure 17, which is the same shape as that in the case where $-ra_2/(a_1 - ra_2) < x < 0$ on (1-1-1). Hence $\mathrm{Area}(S(x)) = V_3$.

Therefore

$$\begin{aligned} s(X,D) &= \mathrm{Vol}(P_{\Sigma,D}) \\ &= \int_{-(r+1)}^{-\frac{a_1}{a_1-ra_2}} \frac{1}{2a_1a_2r^2}\left\{(a_1-2a_2r)x+2a_2r+a_1-a_1r\right\}(x+r+1)dx \\ &\quad + \int_{-\frac{a_1}{a_1-ra_2}}^{-\frac{a_1-a_2}{a_2}} V_5\,dx + \int_{-\frac{a_1-a_2}{a_2}}^{-1} V_1\,dx + \int_{-1}^{-\frac{ra_2}{a_1-ra_2}} V_2\,dx + \int_{-\frac{ra_2}{a_1-ra_2}}^{0} V_3\,dx \\ &= \frac{1}{6a_1a_2^2(a_2r-a_1)}(a_2^3r^3+6a_2^3r^2-3a_1a_2^2r^2+12a_2^3r-12a_1a_2^2r \\ &\quad +3a_1^2a_2r+2a_2^3-12a_1a_2^2+6a_1^2a_2-a_1^3), \end{aligned}$$

(see Section 4 (%o3)).

(1-2-2) Suppose that $r \geq 2$. If $-(a_1-a_2)/a_2 < x < -ra_2/(a_1-ra_2)$, then $S(x)$ is a trapezoid as in the figure 18, which is the same shape as that in the case where $-(a_1-a_2)/a_2 < x < -1$ on (1-2-1). Hence $\mathrm{Area}(S(x)) = V_1$.

If $-ra_2/(a_1-ra_2) < x < -1$, then $S(x)$ is a pentagon as in the figure 19, which is the same shape as that in the case where $-ra_2/(a_1-ra_2) < x < -1$ on (1-1-2). Hence $\mathrm{Area}(S(x)) = V_4$

If $-1 < x < 0$, then $S(x)$ is a pentagonas as in the figure 20, which is the same shape as that in the case where $-ra_2/(a_1-ra_2) < x < 0$ on (1-2-1). Hence $\mathrm{Area}(S(x)) = V_3$.

Therefore

$$\begin{aligned} s(X,D) &= \mathrm{Vol}(P_{\Sigma,D}) \\ &= \int_{-(r+1)}^{-\frac{a_1}{a_1-ra_2}} \frac{1}{2a_1a_2r^2}\left\{(a_1-2a_2r)x+2a_2r+a_1-a_1r\right\}(x+r+1)dx \\ &\quad + \int_{-\frac{a_1}{a_1-ra_2}}^{-\frac{a_1-a_2}{a_2}} V_5\,dx + \int_{-\frac{a_1-a_2}{a_2}}^{-\frac{ra_2}{a_1-ra_2}} V_1\,dx + \int_{-\frac{ra_2}{a_1-ra_2}}^{-1} V_4\,dx + \int_{-1}^{0} V_3\,dx \\ &= \frac{1}{6a_1a_2^2(a_2r-a_1)}(a_2^3r^3+6a_2^3r^2-3a_1a_2^2r^2+12a_2^3r-12a_1a_2^2r \\ &\quad +3a_1^2a_2r+2a_2^3-12a_1a_2^2+6a_1^2a_2-a_1^3), \end{aligned}$$

(see Section 4 (%o4)).

(1-3) We assume that $E_t \leq K_t$. That is, $1/a_2 \leq (r+1)/a_1$. This is equivalent to $a_1 \leq (r+1)a_2$. We also assume that $r \geq 2$. Then $a_1 > ra_2 \geq 2a_2$. Hence $2/a_1 < 1/a_2$ and $(r+2)/a_1 < 2/a_2$. Hence $I_t < G_t$ and $L_t < F_t$. (See the figure 21.)

By the same argument in the case (1-2), $C = L$ and $D = K$ imply $x = -(r+1)$. If $A_t = E_t$, then $-x/a_1 = 1/a_2$. Hence $x = -a_1/a_2$. If

$B = K$, then $(-x/a_1 + 1/a_1, x/r) = ((r+2)/a_1, -1)$. Hence $x = -r$. By the same argument in the case (1-1), we have the following two values of x. If $C = I$ and $D = H$, then $x = -1$. If $A = O$ and $B = H$, then $x = 0$.

Since $ra_2 < a_1 \leq (r+1)a_2$, we have $-(r+1) \leq -a_1/a_2 < -r$. Therefore

$$-(r+1) < -\frac{a_1}{a_2} < -r < -1 < 0.$$

For various values of x, we consider shapes of $S(x)$ and areas of them.

If $-(r+1) < x < -a_1/a_2$, then $S(x)$ is a rectangle as in the figure 22. Hence

$$\begin{aligned} \mathrm{Area}(S(x)) &= \left\{\frac{1}{r}x + \frac{1}{r} - (-1)\right\} \cdot \frac{1}{a_1} \\ &= \frac{1}{a_1 r}(x + r + 1). \end{aligned}$$

We denote this area by V_6.

If $-a_1/a_2 < x < -r$, then $S(x)$ is a pentagon as in the figure 23, which is the same shape as that in the case where $-a_1/(a_1 - ra_2) < x < -(a_1 - a_2)/a_2$ on (1-2). Hence $\mathrm{Area}(S(x)) = V_5$.

If $-r < x < -1$, then $S(x)$ is a pentagon as in the figure 24, which is the same shape as that in the case where $-ra_2/(a_1 - ra_2) < x < -1$ on (1-1-2). Hence $\mathrm{Area}(S(x)) = V_4$.

If $-1 < x < 0$, then $S(x)$ is a pentagon as in the figure 25, which is the same shape as that in the case where $-ra_2/(a_1 - ra_2) < x < 0$ on (1-2-1). Hence $\mathrm{Area}(S(x)) = V_3$.

Therefore

$$\begin{aligned} s(X, D) &= \mathrm{Vol}(P_{\Sigma,D}) \\ &= \int_{-(r+1)}^{-\frac{a_1}{a_2}} V_6\, dx + \int_{-\frac{a_1}{a_2}}^{-r} V_5\, dx + \int_{-r}^{-1} V_4\, dx + \int_{-1}^{0} V_3\, dx \\ &= -\frac{a_2^2 r^2 - 2a_1 a_2 r - 6a_2^2 + a_1^2}{6a_1 a_2^2}, \end{aligned}$$

(see Section 4 (%o5)).

(1-4) We assume that $E_t \leq K_t$. By the same argument in the case (1-3), this is equivalent to $a_1 \leq (r+1)a_2$. We assume that $r = 1$. Then $a_1 \leq 2a_2$, thus $1/a_2 \leq 2/a_1$. Hence $G_t \leq I_t$. We also assume that $3a_2 < 2a_1$. Then $3/a_1 < 2/a_2$. Hence $L_t < F_t$. (See the figure 26.)

We denote the intersection points of the line FG with l_3 by V. Then

$$V = \left(\frac{2-r}{a_1 - ra_2}, -\frac{2a_2 - a_1}{a_1 - ra_2}\right).$$

Since $r = 1$, we have $V = (1/(a_1 - a_2), (a_1 - 2a_2)/(a_1 - a_2))$.

If $C = V$, then we have $(-x/a_1 + 1/a_1, x/r + 1/r) = ((2 - r)/(a_1 - ra_2), -(a_1 - 2a_2)/(a_1 - ra_2))$. Hence $x = -(a_1 + r(a_2 - a_1))/(a_1 - ra_2)$. In particular, $x = -a_2/(a_1 - a_2)$. By the same argument in the case (1-3), we have the following two values of x. If $A_t = E_t$, then $x = -a_1/a_2$. If $B = K$, then $x = -r$, in particular $x = -1$. By the same argument in the case (1-2), we have the following two values of x. If $C = L$ and $D = K$, then $x = -(r + 1)$, in particular, $x = -2$. If $B_t = E_t$, then $x = -(a_1 - a_2)/a_2$. By the same argument in the case (1-1), we have the following two values of x. If $C = I$ and $D = H$, then $x = -1$. If $A = O$ and $B = H$, then $x = 0$.

Since $a_1 \leq 2a_2$, we have $-(r + 1) = -2 \leq -a_1/a_2$. Since $3a_2 < 2a_1$, we have $a_2 < 2(a_1 - a_2)$. Hence $-2 \leq -a_2/(a_1 - a_2)$. If $-a_2/(a_1 - a_2) > -a_1/a_2$, then $a_1^2 - a_2a_1 - a_2^2 > 0$. Therefore $2a_1 < (1 - \sqrt{5})a_2$ or $(1 + \sqrt{5})a_2 < 2a_1$. Since $0 < a_2 < a_1$, we have $2a_1 > (1 + \sqrt{5})a_2$. On the other hand, if $-a_2/(a_1 - a_2) \leq -a_1/a_2$, then $2a_1 \leq (1 + \sqrt{5})a_2$. Since $a_2 < a_1$, we have $-a_1/a_2 < -1$. Since $a_1 \leq 2a_2$, we have $a_1 - a_2 \leq a_2$. Hence $-a_2/(a_1 - a_2) \leq -1 \leq -(a_1 - a_2)/a_2$.

For various values of x, we consider shapes of $S(x)$ and areas of them. (1-4-1) Suppose that $2a_1 > (1 + \sqrt{5})a_2$. Then we have

$$-(r + 1) = -2 < -\frac{a_1}{a_2} < -\frac{a_2}{a_1 - a_2} < -1 < -\frac{a_1 - a_2}{a_2} < 0.$$

If $-(r + 1) = -2 < x < -a_1/a_2$, then $S(x)$ is a rectangle as in the figure 27, which is the same shape as that in the case where $-(r+1) < x < -a_1/a_2$ on (1-3). Hence $\mathrm{Area}(S(x)) = V_6$.

If $-a_1/a_2 < x < -a_2/(a_1 - a_2)$, then $S(x)$ is a pentagon as in the figure 28, which is the same shape as that in the case where $-a_1/(a_1 - ra_2) < x < -(a_1 - a_2)/a_2$ on (1-2-1). Hence $\mathrm{Area}(S(x)) = V_5$.

If $-a_2/(a_1 - a_2) < x < -1$, then $S(x)$ is a hexagon as in the figure 29. Hence

$$\begin{aligned}
\mathrm{Area}(S(x)) &= \frac{1}{a_1}\left(\frac{1}{r}x + \frac{1}{r} + 1\right) - \frac{1}{2}\left(\frac{a_2}{a_1}x + 1\right)\cdot\left(\frac{1}{a_2} + \frac{1}{a_1}\right) \\
&\quad -\frac{1}{2}\left\{-\frac{1}{a_1}x + \frac{1}{a_1} - \left(\frac{1}{a_2} - \frac{1}{ra_2}x - \frac{1}{ra_2}\right)\right\}\cdot\left\{\frac{1}{r}x + \frac{1}{r} - \left(\frac{a_2}{a_1}x - \frac{a_2}{a_1} + 1\right)\right\} \\
&= \frac{1}{ra_1}(x + r + 1) - \frac{1}{2a_1^2a_2}(a_2x + a_1)^2 \\
&\quad -\frac{1}{2a_1^2a_2r^2}\left\{(a_1 - ra_2)x + a_1 + ra_2 - ra_1\right\}^2.
\end{aligned}$$

We denote this area by V_7.

If $-1 < x < -(a_1 - a_2)/a_2$, then $S(x)$ is a hexagon as in the figure 30. Hence

$$\begin{aligned}\mathrm{Area}(S(x)) &= \frac{1}{a_1}\cdot\left(-\frac{1}{r}x\right) - \frac{1}{2}\left(\frac{a_2}{a_1}x - \frac{1}{r}x\right)\cdot\left\{-\frac{1}{ra_2}x - \left(-\frac{1}{a_1}x\right)\right\} \\ &\quad -\frac{1}{2}\left(-\frac{1}{a_1}x + \frac{1}{a_1} - \frac{1}{a_1}\right)\cdot\left\{-\left(\frac{a_2}{a_1}x - \frac{a_2}{a_1} + 1\right)\right\} \\ &= -\frac{1}{ra_1}x - \frac{1}{2a_1^2a_2r^2}(a_1 - ra_2)x^2 - \frac{1}{2a_1^2a_2}(a_2x + a_1 - a_2)^2.\end{aligned}$$

We denote this area by V_8.

If $-(a_1-a_2)/a_2 < x < 0$, then $S(x)$ is a pentagon as in the figure 31, which is the same shape as that in the case where $-ra_2/(a_1 - ra_2) < x < 0$ on (1-1-1). Hence $\mathrm{Area}(S) = V_3$.

Therefore

$$\begin{aligned}s(X, D) &= \mathrm{Vol}(P_{\Sigma,D}) \\ &= \left(\int_{-2}^{-\frac{a_1}{a_2}} V_6\,dx + \int_{-\frac{a_1}{a_2}}^{-\frac{a_2}{a_1-a_2}} V_5\,dx\right. \\ &\quad \left. + \int_{-\frac{a_2}{a_1-a_2}}^{-1} V_7\,dx + \int_{-1}^{-\frac{a_1-a_2}{a_2}} V_8\,dx + \int_{-\frac{a_1-a_2}{a_2}}^{0} V_3\,dx\right)\Bigg|_{r=1} \\ &= \frac{1}{6a_1^2a_2^4(a_2 - a_1)}(2a_2^6 + 4a_1a_2^5 + 2a_2^5 + 3a_1^2a_2^4 - 7a_1a_2^4 - 18a_1^3a_2^3 + 11a_1^2a_2^3 \\ &\quad +15a_1^4a_2^2 - 10a_1^3a_2^2 - 6a_1^5a_2 + 5a_1^4a_2 + a_1^6 - a_1^5),\end{aligned}$$

(see Section 4 (%o7)).

(1-4-2) Suppose that $2a_1 \le (1 + \sqrt{5})a_2$. Then we have

$$-(r+1) = -2 < -\frac{a_2}{a_1 - a_2} < -\frac{a_1}{a_2} < -1 < -\frac{a_1 - a_2}{a_2} < 0.$$

If $-(r + 1) = -2 < x < -a_2/(a_1 - a_2)$, then $S(x)$ is a rectangle as in the figure 32, which is the same shape as that in the case where $-(r + 1) < x < -a_1/a_2$ on (1-3). Hence $\mathrm{Area}(S(x)) = V_6$.

If $-a_2/(a_1 - a_2) < x < -a_1/a_2$, then $S(x)$ is a pentagon as in the figure 33. Hence

$$\begin{aligned}\mathrm{Area}(S(x)) &= \left\{\frac{1}{r}x + \frac{1}{r} - (-1)\right\} \\ &\quad -\frac{1}{2}\left\{\frac{1}{r}x + \frac{1}{r} - \left(\frac{a_2}{a_1}x - \frac{a_2}{a_1} + 1\right)\right\}\cdot\left\{-\frac{1}{a_1}x + \frac{1}{a_1} - \left(\frac{1}{a_2} - \frac{1}{ra_2}x - \frac{1}{ra_2}\right)\right\} \\ &= \frac{1}{a_1r}(x + r + 1) - \frac{1}{2a_1^2a_2r^2}\left\{(a_1 - ra_2)x + a_1 + ra_2 - ra_1\right\}^2.\end{aligned}$$

We denote this area by V_9.

If $-a_1/a_2 < x < -1$, then $S(x)$ is a hexagon as in the figure 34, which is the same shape as that in the case where $-a_2/(a_1 - a_2) < x < -1$ on (1-4-1). Hence $\mathrm{Area}(S(x)) = V_7$.

If $-1 < x < 0$, then $S(x)$ is the same as that in the case where $-1 < x < 0$ on (1-4-1).

Therefore

$$\begin{aligned} s(X,D) &= \mathrm{Vol}(P_{\Sigma,D}) \\ &= \left(\int_{-2}^{-\frac{a_2}{a_1-a_2}} V_6\,dx + \int_{-\frac{a_2}{a_1-a_2}}^{-\frac{a_1}{a_2}} V_9\,dx \right. \\ &\quad \left. + \int_{-\frac{a_1}{a_2}}^{-1} V_7\,dx + \int_{-1}^{-\frac{a_1-a_2}{a_2}} V_8\,dx + \int_{-\frac{a_1-a_2}{a_2}}^{0} V_3\,dx \right)\Bigg|_{r=1} \\ &= \frac{1}{6a_1^2a_2^4(a_2-a_1)}(2a_2^6 + 4a_1a_2^5 + 2a_2^5 + 3a_1^2a_2^4 - 7a_1a_2^4 - 18a_1^3a_2^3 + 11a_1^2a_2^3 \\ &\quad +15a_1^4a_2^2 - 10a_1^3a_2^2 - 6a_1^5a_2 + 5a_1^4a_2 + a_1^6 - a_1^5), \end{aligned}$$

(see Section 4 (%o9)).

(1-5) We assume that $E_t \le K_t$. By the same argument in the case (1-3), we have $a_1 \le (r+1)a_2$. We also assume that $r = 1$. Then $G_t \le I_t$. We also assume that $3a_2 \ge 2a_1$. Then $F_t \le L_t$. (See the figure 35.)

If $B_t = F_t$, then $-x/a_1 + 1/a_1 = 2/a_2$. Hence $x = -(2a_1 - a_2)/a_2$. By the same argument in the case (1-3), we have the following two values of x. If $A_t = E_t$, then $x = -a_1/a_2$. If $B = K$, then $x = -r$, in particular $x = -1$. By the same argument in the case (1-2), we have the following two values of x. If $C = L$ and $D = K$, then $x = -(r+1)$, in particular, $x = -2$. If $B_t = E_t$, then $x = -(a_1 - a_2)/a_2$. By the same argument in the case (1-1), we have the following two values of x. If $C = I$ and $D = H$, then $x = -1$. If $A = O$ and $B = H$, then $x = 0$.

Since $2a_1 \le 3a_2$, we have $2a_1 - a_2 \le 2a_2$. Hence $-2 \le -(2a_1 - a_2)/a_2$. Since $a_1 < a_1 + (a_1 - a_2)$, we have $-(2a_1 - a_2)/a_2 < -a_1/a_2$. Since $a_2 < a_1$, we have $-a_1/a_2 < -1$. Since $a_1 \le 2a_2$, we have $a_1 - a_2 \le a_2$. Hence $-1 \le -(a_1 - a_2)/a_2$. Then we have

$$-(r+1) = -2 < -\frac{2a_1 - a_2}{a_2} < -\frac{a_1}{a_2} < -1 < -\frac{a_1 - a_2}{a_2} < 0.$$

For various values of x, we consider shapes of $S(x)$ and areas of them.

If $-(r+1) = -2 < x < -(2a_1 - a_2)/a_2$, then $S(x)$ is a trapezoid as in the figure 36. Hence

$$\begin{aligned}\mathrm{Area}(S(x)) &= \frac{1}{2}\left\{\left(\frac{1}{a_2}x - \frac{1}{ra_2}x - \frac{1}{ra_2} - \left(-\frac{1}{a_1}x\right)\right)\right\} \cdot \left\{\frac{1}{r}x + \frac{1}{r} - (-1)\right\} \\ &= \frac{1}{2a_1r^2a_2}(x+r+1)\left\{(2a_2r - a_1)x + 3a_1r - a_1\right\}.\end{aligned}$$

If $-(2a_1 - a_2)/a_2 < x < -a_1/a_2$, then $S(x)$ is a pentagon as in the figure 37, which is the same shape as that in case where $-a_2/(a_1 - a_2) < x < -a_1/a_2$ on (1-4-2). Hence $\mathrm{Area}(S(x)) = V_9$.

If $-a_1/a_2 < x < -1$, then $S(x)$ is a hexagon as in the figure 38, which is the same shape as that in case where $-a_2/(a_1 - a_2) < x < -1$ on (1-4-1). Hence $\mathrm{Area}(S(x)) = V_7$.

If $-1 < x < -(a_1 - a_2)/a_2$, then $S(x)$ is a hexagon as in the figure 39, which is the same shape as that in case where $-1 < x < -(a_1 - a_2)/a_2$ on (1-4-1). Hence $\mathrm{Area}(S(x)) = V_8$.

If $-(a_1 - a_2)/a_2 < x < 0$, then $S(x)$ is a pentagon as in the figure 40, which is the same shape as that in case where $-ra_2/(a_1 - ra_2) < x < 0$ on (1-1-1). Hence $\mathrm{Area}(S(x)) = V_3$.

Therefore

$$\begin{aligned}s(X,D) &= \mathrm{Vol}(P_{\Sigma,D}) \\ &= \left(\int_{-2}^{-\frac{2a_1-a_2}{a_2}} \frac{1}{2a_1r^2a_2}(x+r+1)\left\{(2a_2r - a_1)x + 3a_1r - a_1\right\} dx\right. \\ &\quad \left. + \int_{-\frac{2a_1-a_2}{a_2}}^{-\frac{a_1}{a_2}} V_9\, dx + \int_{-\frac{a_1}{a_2}}^{-1} V_7\, dx + \int_{-1}^{-\frac{a_1-a_2}{a_2}} V_8\, dx + \int_{-\frac{a_1-a_2}{a_2}}^{0} V_3\, dx\right)\Bigg|_{r=1} \\ &= \frac{1}{6a_1^2a_2^4}(2a_2^5 - 21a_1a_2^4 + 2a_2^4 + 36a_1^2a_2^3 - 5a_1a_2^3 - 18a_1^3a_2^2 \\ &\quad + 6a_1^2a_2^2 + 5a_1^4a_2 - 4a_1^3a_2 - a_2^5 + a_1^4),\end{aligned}$$

(see Section 4 (%o11)).

(2) Suppose that P_D is a triangle, i.e., $a_1 \le ra_2$. Then the facets of P_D are defined by the equations $x = -a_1$, $y = 0$, and $-x + ry = 0$. Hence we have $I_D = \{1, 3, 4\}$ and

$$P_{\Sigma,D} = \left\{ (x,y,t) \in (M_D \times \mathbb{Z})_{\mathbb{R}} \;\middle|\; \begin{array}{l} 0 \le x + a_1t < 1, \\ 0 \le -y < 1, \\ 0 \le -x + ry < 1 \end{array} \right\}.$$

The point of the intersection of the three hyperplanes defined by $x + a_1t = 0$, $-y = 0$, and $-x + ry = 1$ (resp. $x + a_1t = 0$, $-y = 1$, and

$-x + ry = 0$; $x + a_1 t = 1$, $-y = 0$, and $-x + ry = 0$) is $(-1, 0, 1/a_1)$ (resp. $(-r, -1, r/a_1)$; $(0, 0, 1/a_1)$).

Let $v_1 = {}^t(-1, 0, 1/a_1)$, $v_2 = {}^t(-r, -1, r/a_1)$, and $v_3 = {}^t(0, 0, 1/a_1)$. Since $P_{\Sigma,D}$ is spanned by v_1, v_2, and v_3, we have

$$s(X, D) = \mathrm{Vol}(P_{\Sigma,D}) = |\det(v_1, v_2, v_3)| = \frac{1}{a_1}.$$

(3) Suppose that P_D is not full-dimensional. P_D is not full-dimensional if and only if $a_1 = 0$ or $a_2 = 0$. First we consider the case where $a_2 = 0$. Then P_D is a line segment whose edges are defined by $x = -a_1$ and $-x + ry = 0$. Hence we have $I_D = \{1, 4\}$. We see that $M_D = \mathbb{Z} \times \{0\} \subset M$ with basis $\{\langle 1, 0\rangle\}$. Let $\pi : N \to N_D = \mathbb{Z} \times \{0\}$ be the first projection map. Since $v_1 \cdot \langle 1, 0\rangle = 1$ (resp. $v_4 \cdot \langle 1, 0\rangle = -1$), $\pi(v_1)$ (resp. $\pi(v_4)$) is N_D-primitive and $c_1 = 1$ (resp. $c_4 = 1$). Therefore we have

$$P'_{\Sigma,D} = \left\{ (x, t) \in (M_D \times \mathbb{Z})_{\mathbb{R}} \;\middle|\; \begin{array}{l} 0 \le x + a_1 t < 1, \\ 0 \le -x < 1 \end{array} \right\}.$$

$P'_{\Sigma,D}$ is the parallelogram spanned by the two vectors $v_1 = {}^t(0, 1/a_1)$ and $v_2 = {}^t(-1, 1/a_1)$. Hence we have

$$s(X, D) = \mathrm{Vol}(P'_{\Sigma,D}) = |\det(v_1, v_2)| = \frac{1}{a_1}.$$

Next we consider the case where $a_1 = 0$. Since $\mathrm{Sec}(X, a_2 D_2) \cong k[x]$, the F-signature is not defined. (An F-signature is defined for the section ring whose dimension is at least two.)

3. Figures

In this section, we gather together the figures in the computation of F-signatures of Hirzebruch surfaces.



FIGURE 1.



FIGURE 2.



FIGURE 3.



FIGURE 4.



FIGURE 5.



FIGURE 6.



FIGURE 7.



FIGURE 8.



FIGURE 9.



FIGURE 10.



FIGURE 11.



FIGURE 12.



FIGURE 13.



FIGURE 14.



FIGURE 15.



FIGURE 16.



FIGURE 17.



FIGURE 18.



FIGURE 19.



FIGURE 20.



FIGURE 21.



FIGURE 22.



FIGURE 23.



FIGURE 24.



FIGURE 25.



FIGURE 26.



FIGURE 27.



FIGURE 28.



FIGURE 29.



FIGURE 30.



FIGURE 31.



FIGURE 32.



FIGURE 33.



FIGURE 34.



FIGURE 35.



FIGURE 36.



FIGURE 37.



FIGURE 38.



FIGURE 39.



FIGURE 40.

4. Maxima source codes

To compute integrals, we use the Maxima which is a computer algebra system. In particular, we use a software for Maxima called wxMaxima. In the following code, we use a (resp. b) instead of a_1 (resp. a_2) The following are source code:

```
(%i1)/*case 1 (2a_2\leq a_1)*/
factor(integrate((1/(2*a^2*b*r^2))*((a-b*r)*x+a+b*r)^2,
x,(b*r+a)/(b*r-a),a/(b*r-a))
+integrate((1/a)*(1/r-b/a)*x+(1/(2*a))*(2/r+b/a),x,a/(b*r-a),-1)
+integrate((b/(2*a^2))*(1-2*x),x,-1,(b*r)/(b*r-a))
+integrate(-(x/(a*r))-(1/2)*(b/a-1/r)*(1/a-1/(b*r))*x^2,
x,(b*r)/(b*r-a),0));
```

(%o1)
$$-\frac{b}{a\ (b\,r-a)}$$

```
(%i2)/*case 1 (2a_2> a_1)*/
factor(integrate(1/(2*a^2*b*r^2)*((a-r*b)*x+a+r*b)^2,
x,-(a+r*b)/(a-b*r),-a/(a-r*b))
+integrate(1/(2*r*a^2)*(2*(a-r*b)*x+2*a+r*b),
x,-a/(a-r*b),-(r*b)/(a-r*b))
+integrate(1/(r*a)-(r*b-a)^2/(2*r^2*a^2*b)*x^2,
x,-(r*b)/(a-r*b),-1)
+integrate(-x/(r*a)-(r*b-a)^2*x^2/(2*r^2*a^2*b),x,-1,0));
```

(%o2)
$$-\frac{b}{a\ (b\,r-a)}$$

```
(%i3)/*case_2 r=1 */
factor(integrate(1/(2*a*b*r^2)*((a-2*b*r)*x+2*b*r+a-a*r)*(x+r+1),
x,-(r+1),-a/(a-r*b))
+integrate((x+r+1)/(r*a)-(b*x+a)^2/(2*a^2*b),x,-a/(a-r*b),-(a-b)/b)
+integrate(1/(2*a^2*r)*(2*(a-r*b)*x+2*a+r*b),x,-(a-b)/b,-1)
+integrate(b*(1-2*x)/(2*a^2),x,-1,-(r*b)/(a-r*b))
+integrate(-x/(r*a)-(r*b-a)^2*x^2/(2*r^2*a^2*b),x,-r*b/(a-r*b),0));
```

(%o3)
$$\frac{b^3\,r^3+6\,b^3\,r^2-3\,a\,b^2\,r^2+12\,b^3\,r-12\,a\,b^2\,r+3\,a^2\,b\,r+2\,b^3-12\,a\,b^2+6\,a^2\,b-a^3}{6\,a\,b^2\ (b\,r-a)}$$

```
(%i4)/*case_2 r>1 */
factor(integrate(1/(2*a*b*r^2)*((a-2*b*r)*x+2*b*r+a-a*r)*(x+r+1),
```

```
x,-(r+1),-a/(a-r*b))
+integrate((x+r+1)/(r*a)-(b*x+a)^2/(2*a^2*b),x,-a/(a-r*b),-(a-b)/b)
+integrate(1/(2*a^2*r)*(2*(a-r*b)*x+2*a+r*b),x,-(a-b)/b,-r*b/(a-r*b))
+integrate(1/(r*a)-(r*b-a)^2*x^2/(2*r^2*a^2*b),x,-(r*b)/(a-r*b),-1)
+integrate(-x/(r*a)-(r*b-a)^2*x^2/(2*r^2*a^2*b),x,-1,0));
```

(%o4)

$$\frac{b^3\,r^3+6\,b^3\,r^2-3\,a\,b^2\,r^2+12\,b^3\,r-12\,a\,b^2\,r+3\,a^2\,b\,r+2\,b^3-12\,a\,b^2+6\,a^2\,b-a^3}{6\,a\,b^2\,\left(b\,r-a\right)}$$

```
(%i5)/*case3 */
factor(integrate((x+r+1)/(a*r),x,-(r+1),-a/b)
+integrate((x+r+1)/(r*a)-(b*x+a)^2/(2*a^2*b),x,-a/b,-r)
+integrate(1/(r*a)-(r*b-a)^2*x^2/(2*r^2*a^2*b),x,-r,-1)
+integrate(-x/(r*a)-(r*b-a)^2*x^2/(2*r^2*a^2*b),x,-1,0));
```

(%o5)

$$-\frac{b^2\,r^2-2\,a\,b\,r-6\,b^2+a^2}{6\,a\,b^2}$$

```
(%i6)/* case4 2a_1>(1+\sqrt{5})*a_2 */
f(r):=integrate((x+r+1)/(a*r),x,-2,-a/b)
+integrate((x+r+1)/(r*a)-(b*x+a)^2/(2*a^2*b),x,-a/b,-b/(a-b))
+integrate((x+r+1)/(r*a)-(b*x+a)^2/(2*a^2*b)
-((a-r*b)*x+a+r*b-r*a)^2/(2*a^2*b*r^2),x,-b/(a-b),-1)
+integrate(-x/(r*a)-(a-r*b)*x^2/(2*a^2*b*r^2)-(b*x+a-b)^2/(2*a^2*b),
x,-1,-(a-b)/b)
+integrate(-x/(r*a)-(r*b-a)^2*x^2/(2*r^2*a^2*b),x,-(a-b)/b,0);
(%i7)factor(f(1));
```

(%o7)

$$\frac{2\,b^6+4\,a\,b^5+2\,b^5+3\,a^2\,b^4-7\,a\,b^4-18\,a^3\,b^3+11\,a^2\,b^3+15\,a^4\,b^2-10\,a^3\,b^2-6\,a^5\,b+5\,a^4\,b+a^6-a^5}{6\,a^2\,b^4\,\left(b-a\right)}$$

```
(%i8)/* case4 2a_1\leq(1+\sqrt{5})*a_2 */
g(r):=integrate((x+r+1)/(a*r),x,-2,-b/(a-b))
+integrate((x+r+1)/(r*a)-((a-r*b)*x+a+r*b-r*a)^2/(2*a^2*b*r^2),
x,-b/(a-b),-a/b)
+integrate((x+r+1)/(r*a)-(b*x+a)^2/(2*a^2*b)
-((a-r*b)*x+a+r*b-r*a)^2/(2*a^2*b*r^2),x,-a/b,-1)
+integrate(-x/(r*a)-(a-r*b)*x^2/(2*a^2*b*r^2)-(b*x+a-b)^2/(2*a^2*b),
x,-1,-(a-b)/b)
+integrate(-x/(r*a)-(r*b-a)^2*x^2/(2*r^2*a^2*b),x,-(a-b)/b,0);
(%i9)factor(g(1));
```

(%o9)
$$\frac{2\,b^6+4\,a\,b^5+2\,b^5+3\,a^2\,b^4-7\,a\,b^4-18\,a^3\,b^3+11\,a^2\,b^3+15\,a^4\,b^2-10\,a^3\,b^2-6\,a^5\,b+5\,a^4\,b+a^6-a^5}{6\,a^2\,b^4\,(b-a)}$$

```
(%i10) /*case 5*/
h(r):=integrate((x+r+1)*((2*b*r-a)*x+3*a*r-a)/(2*a*r^2*b),
x,-2,-(2*a-b)/b)
+integrate((x+r+1)/(r*a)-((a-r*b)*x+a+r*b-r*a)^2/(2*a^2*b*r^2),
x,-(2*a-b)/b,-a/b)
+integrate((x+r+1)/(r*a)-(b*x+a)^2/(2*a^2*b)
-((a-r*b)*x+a+r*b-r*a)^2/(2*a^2*b*r^2),x,-a/b,-1)
+integrate(-x/(r*a)-(a-r*b)*x^2/(2*a^2*b*r^2)-(b*x+a-b)^2/(2*a^2*b),
x,-1,-(a-b)/b)
+integrate(-x/(r*a)-((r*b-a)*x)^2/(2*r^2*a^2*b),x,-(a-b)/b,0);
(%i11)factor(h(1));
```

(%o11)
$$\frac{2\,b^5-21\,a\,b^4+2\,b^4+36\,a^2\,b^3-5\,a\,b^3-18\,a^3\,b^2+6\,a^2\,b^2+5\,a^4\,b-4\,a^3\,b-a^5+a^4}{6\,a^2\,b^4}$$

References

[CLS11] David A. Cox, John B. Little, and Henry K. Schenck, *Toric varieties*, Graduate Studies in Mathematics, vol. 124, American Mathematical Society, Providence, RI, 2011. MR 2810322

[HN15] Mitsuyasu Hashimoto and Yusuke Nakajima, *Generalized F-signature of invariant subrings*, J. Algebra **443** (2015), 142–152. MR 3400399

[HL02] Craig Huneke and Graham J. Leuschke, *Two theorems about maximal Cohen-Macaulay modules*, Math. Ann. **324** (2002), no. 2, 391–404. MR 1933863

[San15] Akiyoshi Sannai, *On dual F-signature*, Int. Math. Res. Not. IMRN (2015), no. 1, 197–211. MR 3340299

[Tuc12] Kevin Tucker, *F-signature exists*, Invent. Math. **190** (2012), no. 3, 743–765. MR 2995185

[Kor12] Michael R. Von Korff, *The F-Signature of Toric Varieties*, ProQuest LLC, Ann Arbor, MI, 2012, Thesis (Ph.D.)–University of Michigan. MR 3093997

Department of General education, National Institute of Technology, Fukushima College, 30 Aza-Nagao, Kamiarakawa, Iwaki-shi, Fukushima 970-8034, Japan

E-mail address: hirose@fukushima-nct.ac.jp

Department of General Education, National Institute of Technology, Fukushima College, Fukushima 970-8034, Japan.

E-mail address: sawada@fukushima-nct.ac.jp